


## CASE STUDY IN HONDURAS: QUEZUNGUAL SLASH AND MULCH AGROFORESTRY SYSTEM

In the early 1990s, FAO began working with Honduran farmers on the steep hillsides of South Western Honduras to mitigate the effects of deforestation and land degradation, and to develop and disseminate new and more sustainable agricultural practices. Farmers there have practiced 'slash and burn' cultivation and abandoned the age-old practice of allowing cleared fields to lie fallow long enough for tree cover to grow back and for the soil to recover. Without trees to anchor the depleted soil, erosion increased which reduced the quality of water and its availability to downstream users. As agricultural production declined, the rates of rural poverty and malnutrition rose sharply. Recognizing the urgent need to change their cultivation practices, Honduran farmers developed a low-cost, resource-conserving system for growing their crops. Instead

of clearing the forests and burning vegetation, they adopted a slash and mulch approach. First, they broadcasted beans and sorghum in an area of well-developed naturally regenerated secondary forest. After planting, they selectively cut and pruned the trees and shrubs and spread the leaves and small branches on the soil surface to create a layer of mulch. High-value timber, fruit and fuelwood trees

were left to grow. Once the beans and sorghum had been harvested, maize was planted. Farmers continued to prune trees to allow sufficient sunlight to reach the crops while leaves, branches and crop residues were used to maintain a semi-permanent soil cover. Honduran farmers have embraced this system as it is founded on familiar, indigenous farming practices and delivers many benefits. By retaining soil moisture and preventing erosion, QSMAS has made farms more resilient to extreme weather events, such as Hurricane Mitch in 1998. The system also reduces the time required to prepare the land and control weeds.

Source: Adapted from *Save and Grow in practice: maize, rice and wheat a guide to sustainable cereal production* (FAO 2016).



COMMON BEANS  
(PHASEOLUS VULGARIS)

### SOURCES:

<sup>1</sup> Nulik J, Dalgliesh N, Cox K and Gabb S. (2013) Integrating herbaceous legumes into crop and livestock systems in eastern Indonesia. Australian Centre for International Agricultural Research (ACIAR), Canberra.

<sup>2</sup> Rose TJ, Hardiputra B, Rengel Z. (2010) Wheat, canola and grain legume access to soil phosphorus fractions differs in soils with contrasting phosphorus dynamics. *Plant and Soil* 326: 159–170.

<sup>3</sup> Blanchart E, Villenave C, Viallatoux A, Barthès B, Girardin C, Azontonde A and Feller C. (2005) Long-term effect of a legume cover crop (*Mucuna pruriens* var. utilis) on the communities of soil macrofauna and nematofauna under maize cultivation, in southern Benin. *European Journal of Soil Biology* 42: 136–144.

<sup>4</sup> Brussaard L, Ruijter PC de and Brwon GG. (2007) Soil biodiversity for agriculture sustainability. *Agriculture, Ecosystems and Environment* 121: 233–244.

<sup>5</sup> Giller KE, Wilson KJ. (1991) Nitrogen fixation in tropical cropping systems. CAB International, Wallingford.

<sup>6</sup> Jensen, ES; Peoples, MB; Boddey, RM; Gresshoff, PM; Hauggaard-Nielsen, H;

Alves, BJR; Morrison MJ. 2012. Legumes for mitigation of climate change and the provision of feedstock for biofuels and biorefineries. A review. *Agronomy for Sustainable Development* 32:329–364.

<sup>7</sup> Atangana A, Khasa D, Chang S, Degrande A (2014). *Tropical Agroforestry*. Springer, Dornrecht.

<sup>8</sup> FAO (2016). *Save and grow in practice – maize, rice and wheat: A guide to a sustainable cereal production*. Food and Agriculture Organization of the United Nations, Rome.

<sup>9</sup> Heller J, Begemann F, Mushonga J, (1997). Bambara groundnut (*Vigna subterranea* (L.) Verdc.): Promoting the conservation and use of underutilized and neglected crops. Institute of Plant Genetics and Crop Plant Research, Gatersleben; Department of Research and Specialist Services, Harare; International Plant Genetic Resources Institute, Rome.

<sup>10</sup> Young A (1991). *Agroforestry for soil conservation*. CAB International, Wallingford.

<sup>11</sup> ICRISAT (2016). *Agroforestry systems*. International Crops Research Institute for the semi-arid Tropics, Hyderabad



Food and Agriculture  
Organization of the  
United Nations




# PULSES AND BIODIVERSITY

**PULSES ARE ABLE TO INCREASE BIODIVERSITY AS THEY ARE ABLE TO FIX THEIR OWN NITROGEN INTO THE SOIL, WHICH INCREASES SOIL FERTILITY.**

It is estimated that there are hundreds of varieties of pulses, including many local varieties that are not exported or grown worldwide. The genetic diversity of these crops is an essential component for on-farm soil and pest management, especially for small-scale farmers.

An important attribute of pulses is their ability to biologically fix nitrogen. These plants, in symbiosis with certain types of bacteria, namely *Rhizobium* and *Bradyrhizobium*, are able to convert atmospheric nitrogen into nitrogen compounds that can be used by plants, while also improving soil fertility.<sup>1</sup>

Some varieties of pulses are also able to free soil-bound phosphorous,



and phosphorous also plays an important role in the nutrition of plants.<sup>2</sup> The presence of pulses in agro-ecosystems helps to maintain

and/or increase vital microbial biomass and activity in the soil, so nourishing the development of those organisms that are responsible for promoting soil structure and nutrient availability.<sup>3</sup>

A high soil biodiversity provides ecosystems with not only greater resistance and resilience against disturbance and stress, but also improves the ability of ecosystems to suppress diseases.<sup>4</sup> All these features are particularly important for mainstreaming soil health, which is the foundation of food security and health.

## KEY FACTS

➤ Good soil health is the foundation of food security and pulses **help** to increase soil **microbial biomass** and activity, thus improving soil biodiversity.

➤ Cultivating pulses in **multiple cropping systems** enriches agro-biodiversity, ensures resilience to climate change and **improves ecosystem services**.

➤ Pulses have a multiple role in promoting living organisms and ecological complexity to **re-establish** the natural **good functioning of ecosystems**.



#IYP2016  
fao.org/pulses-2016

© FAO 2016  
15389E/1/02.16



# MULTIPLE CROPPING SYSTEMS AND BIODIVERSITY

Pulses cannot improve on-farm diversity *per se*. This means that if a farmer changes from cultivating only cereal species to cultivating only pulse species, on-farm diversity does not change. That said, pulses are a crucial component of multiple cropping systems, namely intercropping, crop rotation and agroforestry. These cropping systems have a higher species diversity than monocrop systems. Increasing species diversity of cropping systems could translate into not only a more efficient use of resources, namely light, water and nutrients<sup>5</sup>, but also into higher outputs as yields are increased, and a lower risk of overall crop failure. The selection of which multiple cropping system to use is less important because the choice will be determined by the individual attributes of each agro-ecosystem.



What is clear is that pulses *should* be an integral part of agro-ecosystems because they bring the agro-systems into balance. Additionally, cropping systems that are more resilient to climate change can be developed through the inclusion of local varieties, such as bambara beans, which are currently not well known or produced.

In multiple cropping systems, services such as nutrient recycling and soil formation are improved through the pulses’ abilities to fix nitrogen and free phosphorous and their capacity to increase soil biodiversity. At the same time, when used in multiple cropping systems, pulses also help to curb and control pests and diseases. Additionally, since pulses often promote higher rates of accumulation of soil carbon than cereals or grasses<sup>6</sup>, they can contribute to improve the carbon sequestration of agro-ecosystems.

Being **drought-tolerant**, **pigeon peas** (*Cajanus cajan* (L.) Huth) are often intercropped with cereals in smallholder farming systems in Asia, Africa and the Caribbean. As pigeon peas are also deep rooting, they do not compete with maize for water.

## BIODIVERSITY BENEFITS OF PULSES IN MULTIPLE CROPPING SYSTEMS

